

8/27/05 10/773,414

NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 17:27:01 ON 27 AUG 2005

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 17:27:12 ON 27 AUG 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 26 AUG 2005 HIGHEST RN 861902-61-6

DICTIONARY FILE UPDATES: 26 AUG 2005 HIGHEST RN 861902-61-6

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JULY 14, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.

\*\*\*\*\*  
\*  
\* The CA roles and document type information have been removed from \*  
\* the IDE default display format and the ED field has been added, \*  
\* effective March 20, 2005. A new display format, IDERL, is now \*  
\* available and contains the CA role and document type information. \*  
\*  
\*\*\*\*\*


Structure search iteration limits have been increased. See HELP SLIMITS for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:

<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10773414a.str



chain nodes :

27 28

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23  
24 25 26

chain bonds :

3-28 6-9 10-18 13-28 14-27

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 13-14 13-17  
14-15 15-16 15-23 15-26 16-17 18-19 18-22 19-20 20-21 21-22 23-24 24-25  
25-26

exact/norm bonds :

13-14 13-17 13-28 14-15 14-27 15-16 15-23 15-26 16-17 18-19 18-22 19-20  
20-21 21-22 23-24 24-25 25-26

exact bonds :

3-28 6-9 10-18

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom  
20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom 27:CLASS 28:CLASS

L1 STRUCTURE UPLOADED

=> d

L1 HAS NO ANSWERS

L1 STR

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

Structure attributes must be viewed using STN Express query preparation.

=> s L1

SAMPLE SEARCH INITIATED 17:27:35 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 55 TO ITERATE

100.0% PROCESSED 55 ITERATIONS  
SEARCH TIME: 00.00.01

2 ANSWERS

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 656 TO 1544  
PROJECTED ANSWERS: 2 TO 124

L2 2 SEA SSS SAM L1

=> s L1 full

FULL SEARCH INITIATED 17:27:41 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 1139 TO ITERATE

100.0% PROCESSED 1139 ITERATIONS  
SEARCH TIME: 00.00.01

42 ANSWERS

L3 42 SEA SSS FUL L1

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION

FULL ESTIMATED COST

161.33

161.54

FILE 'CAPLUS' ENTERED AT 17:27:44 ON 27 AUG 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 27 Aug 2005 VOL 143 ISS 10

FILE LAST UPDATED: 26 Aug 2005 (20050826/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s L3

L4 677 L3

=> s L4 and (synthesi? or "process of making" or "method of making")  
1439123 SYNTHESI?

```

2135433 "PROCESS"
1428941 "PROCESSES"
3177531 "PROCESS"
      ("PROCESS" OR "PROCESSES")
      0 "OF"
      195 "OFS"
      195 "OF"
      ("OF" OR "OFS")
257447 "MAKING"
      31 "MAKINGS"
257472 "MAKING"
      ("MAKING" OR "MAKINGS")
      0 "PROCESS OF MAKING"
      ("PROCESS" (W) "OF" (W) "MAKING")
2918309 "METHOD"
1204277 "METHODS"
3783802 "METHOD"
      ("METHOD" OR "METHODS")
      0 "OF"
      195 "OFS"
      195 "OF"
      ("OF" OR "OFS")
257447 "MAKING"
      31 "MAKINGS"
257472 "MAKING"
      ("MAKING" OR "MAKINGS")
      0 "METHOD OF MAKING"
      ("METHOD" (W) "OF" (W) "MAKING")
L5      36 L4 AND (SYNTHESI? OR "PROCESS OF MAKING" OR "METHOD OF MAKING")
=> d ibib abs fhitr 1-36

```

ACCESSION NUMBER: 2005:732585 CAPLUS

DOCUMENT NUMBER: 143:179169

TITLE: Cosmetic compositions ACE inhibitors and/or angiotensin II receptor antagonists for treatment of skin aging

INVENTOR(S): Jensen, Benny Vittrup

PATENT ASSIGNEE(S): Ace Aps, Den.

SOURCE: PCT Int. Appl., 48 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005072696	A1	20050811	WO 2005-DK65	20050128
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	EW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

PRIORITY APPLN. INFO.: DK 2004-136 A 20040130  
US 2004-553661P P 20040316

AB The present invention relates to a method and cosmetic preparation

comprising

an ACE inhibitor and/or angiotensin II receptor antagonist present in an amount of about 0.01 to 100 mg/kg each for the treatment of skin aging or wrinkling. For example, an ACE inhibitor, such as lisinopril 10 mg/kg was formulated in a cream base comprising (i) Phase A containing Emulgade SE

4.0%.

Cutina MD 1.0%, Lanette O 1.0%, Baysilon M 350 0.5%, Cetiol PGL 7.0%, Cetiol OE 4.0%, and Copherol 1250 0.5%, (ii) Phase B containing D-panthenol 1.0%, glycerin (86%) 5.0%, and water 71.5%, (iii) Phase C containing

Carbopol

980 0.2% and Cetiol PGL 1.0%, and (iv) Phase C containing KDH (20%) 0.3% and perfums/preservative as needed.

IT

138402-11-6, Avapro  
RL: COS (Cosmetic use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(Irbesartan; compns. containing ACE inhibitors and/or angiotensin II receptor antagonists for improvement and maintenance of skin tone and treatment of skin aging)

RN

138402-11-6 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)

ACCESSION NUMBER: 2005:640217 CAPLUS

TITLE: Improved synthesis of irbesartan, an antihypertensive active pharmaceutical ingredient

AUTHOR(S): Sumalatha, Bollikonda Satyanarayana Yasareni;

Venkatraman, Sundram; Reddy, Ghanta Mahesh; Reddy, Padi Pratap

CORPORATE SOURCE: Research and Development Centre, Dr Reddy's

Laboratories Limited, Hyderabad, India

SOURCE: Synthetic Communications (2005), 35(14), 1979-1982

CODEN: SYNCAY; ISSN: 0039-7911

PUBLISHER: Taylor &amp; Francis, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An improved synthesis of the antihypertensive drug irbesartan I, based on the Suzuki reaction, was described.

IT

INDEXING IN PROGRESS

IT 138402-11-6P, Irbesartan

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation of irbesartan antihypertensive active pharmaceutical

ingredient


based on Suzuki reaction)

RN 138402-11-6 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2005:612299 CAPLUS

DOCUMENT NUMBER: 143:133380

TITLE: Preparation of azabicyclic heterocycles as cannabinoid receptor modulators

INVENTOR(S): Gu, Guixue; Ewing, William R.; Mikkilineni, Amarendra

B.; Fendri, Annapurna; Ellsworth, Bruce A.; Sher, Philip M.; Gerritz, Samuel; Sun, Chongqing; Murugesan, Natesan; Wu, Ximao

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 101 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005063762	A1	20050714	WO 2004-US42878	20041217
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	EW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2005171110	A1	20050804	US 2004-16198	20041217
PRIORITY APPLN. INFO.:			US 2003-531451P	P 20031219
			US 2004-16198	A 20041217

GI




I



II

AB The present application describes compds. I (R1, R2 = halo, CN, alkyl, etc.; R3 = H alkyl, alkenyl, cycloalkyl, etc.; R4 is absent when n is a double bond; R4 = H, alkyl, cycloalkyl, etc.; R5 = halo, (un)substituted OH, NH2, etc. when n is a single bond; R5 = O when n = a double bond; n = a single or double bond; when n is a single bond, n is a double bond; when n is a double bond, n is a single bond), pharmaceutical compns. comprising at least one compound I and optionally one or more addnl. therapeutic agents and methods of treatment using the compds. I both alone and in combination with one or more addnl. therapeutic agents. Over 40

L5 ANSWER 3 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 compds. I were prepd. E.g., a multi-step synthesis of II, starting from dichloromandelic anhydride, was given. The exemplified compds. I showed the CB-1 receptor binding Ki values in the range of 0.01 nM to 10000 nM.  
 IT 138402-11-6  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (co-drug; preparation of azabicyclic heterocycles as cannabinoid receptor modulators)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L5 ANSWER 4 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2005:572592 CAPLUS  
 DOCUMENT NUMBER: 143:97378  
 TITLE: Preparation of azabicyclic heterocycles as cannabinoid receptor modulators  
 INVENTOR(S): Yu, Guixue; Ewing, William R.; Mikkilineni, Anarendra B.; Pendri, Annapurna Sher, Philip M.; Gerritz, Samuel; Ellsworth, Bruce A.; Wu, Gang; Huang, Yanting; Sun, Chongqing; Murugesan, Natesan; Gu, Zhengxiang; Wang, Ying; Sitkoff, Doree; Johnson, Stephen R.; Wu, Ximao  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 196 pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005143381	A1	20050630	US 2004-16135	20041217
WO 2005063761	A1	20050714	WO 2004-0542820	20041217
V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZH, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, BJ, CF, CG, CI, CN, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2005061509	A1	20050707	WO 2004-0542542	20041220
V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZH, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, BJ, CF, CG, CI, CN, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: US 2003-531451P P 20031219  
 US 2004-16135 A 20041217  
 GI




L5 ANSWER 4 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 AB The present application describes compds. I [R1, R2 = halo, CN, alkyl, etc.; R3 = alkyl, alkenyl, cycloalkyl, etc.; R6 = H, alkyl, cycloalkyl, etc.; R7 is absent when double bonds; or R7 = H, alkyl, cycloalkyl, etc.], pharmaceutical compns. comprising at least one compound I and optionally one or more addnl. therapeutic agents and methods of treatment using the compds. I both alone and in combination with one or more addnl. therapeutic agents. Over 400 compds. I were prepared E.g., a multi-step synthesis of II, starting from dibromopyridazinone, was given. Representative compds. I showed the CB-1 receptor binding Ki values in the range of 0.01 nM to 10000 nM.  
 IT 138402-11-6, Irbesartan  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (co-drug; preparation of azabicyclic heterocycles as cannabinoid receptor modulators)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 5 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2005:474941 CAPLUS  
 DOCUMENT NUMBER: 142:476245  
 TITLE: Synergistic effect of amlodipine and atorvastatin on aortic endothelial cell nitric oxide release, and therapeutic use  
 INVENTOR(S): Mason, R. Preston  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 19 pp., Cont.-in-part of U.S. Ser. No. 921,479.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005119270	A1	20050602	US 2004-987271	20041112
US 2002052394	A1	20020502	US 2001-921479	20010803
US 6835742	B2	20041228		
US 2005009888	A1	20050113	US 2004-911807	20040805
PRIORITY APPLN. INFO.: US 2000-223214P P 20000804 US 2001-921479 A2 20010803				


AB The combination of amlodipine and atorvastatin acts to synergistically synthesize NO production. Moreover, the addition of a tertiary compound complements this combination of amlodipine and atorvastatin in NO production. The combination of the invention may be used to treat arterial and related heart disease.  
 IT 138402-11-6, Irbesartan  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (synergistic effect of amlodipine and atorvastatin on aortic endothelial cell nitric oxide release, and therapeutic use)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 6 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
ACCESSION NUMBER: 2005:371095 CAPLUS  
DOCUMENT NUMBER: 142:423895  
TITLE: Methods for controlling mast cell-derived renin and  
uses in treating conditions with abnormal renin levels  
INVENTOR(S): Silver, Randi B.; Levi, Roberto  
PATENT ASSIGNEE(S): Cornell Research Foundation, Inc., USA  
SOURCE: PCT Int. Appl., 129 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005037317	A2	20050428	WO 2004-US33755	20041013
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			



PRIORITY APPL. INFO.: US 2003-512142P P 20031017  
AB The invention relates to the discovery that renin is present in mast cells and can act in a localized manner to initiate and/or exacerbate a number of conditions. Thus, the invention provides methods for treating cardiac, vascular, lung, liver, cervical, intestinal, muscle, pancreatic, brain, kidney, skin and other conditions that involve inhibiting the synthesis and/or release of renin from mast cells and/or inhibiting the activity of renin after release from mast cells. The methods of the invention can also involve inhibiting elements of the local renin-angiotensin system (e.g. inhibiting ACE and angiotensin II receptors), thereby inhibiting angiotensin II produced locally from mast-cell-derived renin.  
IT 138402-11-6, Irbesartan  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (methods for controlling mast cell-derived renin and uses in treating conditions with abnormal renin levels)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 7 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
ACCESSION NUMBER: 2004:1127349 CAPLUS  
DOCUMENT NUMBER: 142:74574  
TITLE: Preparation of 1,2,4-triazolyethylamines as  
modulators of the glucocorticoid receptor  
INVENTOR(S): Robinson, Leslie; Rueter, Jaimie K.; Morse, Wilna J.  
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
SOURCE: PCT Int. Appl., 69 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004111015	A1	20041223	WO 2004-US18487	20040611
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

US 2004266831 A1 20041230 US 2004-865443 20040610  
PRIORITY APPL. INFO.: US 2003-477545P P 20030611  
OTHER SOURCE(S): MARPAT 142:74574  
GI



AB Title compds. I [A, B = cycloalkyl, aryl, heteroaryl; R1 = H, acyl, carbonyl, etc.; R2-4 = H, alkyl, heteroalkyl, etc.; R5-6 = H, F, Cl, Br.

L5 ANSWER 7 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
etc.] are prepd. General synthetic procedures are provided for the synthesis of 19 examples, e.g., II. Example compds. are tested in a glucocorticoid receptor binding assay in the range of 0.1 nM to 40 µM [no data]. I are glucocorticoid receptor modulators and are useful in treating diseases requiring glucocorticoid receptor agonist or antagonist therapy such as obesity, diabetes, inflammatory and immune disorders.  
IT 138402-11-6, Irbesartan  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (combination pharmaceutical preparation of 1,2,4-triazolyethylamines as modulators of glucocorticoid receptor)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 8 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
ACCESSION NUMBER: 2004:1124587 CAPLUS  
DOCUMENT NUMBER: 142:69188  
TITLE: Combination therapy for the treatment of diabetes  
INVENTOR(S): Erondou, Ngozi E.; Fong, Tung M.; MacNeil, Douglas J.; Van Der Ploeg, Leonardus H. T.; Kanatani, Akio  
PATENT ASSIGNEE(S): Merck & Co., Inc., USA; Banyu Pharmaceutical Co., Ltd.  
SOURCE: PCT Int. Appl., 109 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004110375	A2	20041223	WO 2004-US17291	20040602
WO 2004110375	A3	20050512		
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BV, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			


PRIORITY APPL. INFO.: US 2003-476388P P 20030606  
OTHER SOURCE(S): MARPAT 142:69188  
AB The present invention relates to compns. comprising an anti-obesity agent and an anti-diabetic agent useful for the treatment of diabetes, diabetes associated with obesity and diabetes-related disorders. The present invention further relates to methods of treating or preventing obesity, and obesity-related disorders, in a subject in need thereof by administering a composition of the present invention. The present invention further provides for pharmaceutical compns., medicaments, and kits useful in carrying out these methods.  
IT 138402-11-6, Irbesartan  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (combination therapy of diabetes and diabetes-related disorders using antiobesity agent and antidiabetic agent and other agents)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 9 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:1086422 CAPLUS  
DOCUMENT NUMBER: 142:197940  
TITLE: Dual Angiotensin II and Endothelin A Receptor Antagonists: Synthesis of 2'-Substituted N-3-Isoxazolyl Biphenylsulfonamides with Improved Potency and Pharmacokinetics  
AUTHOR(S): Murugesan, Natesan; Gu, Zhengxiang; Fadnis, Leena; Talley, John E.; Baska, Rose Ann F.; Yang, Yifan; Beyer, Sophie M.; Monshizadegan, Hossain; Dickinson, Kenneth E.; Valentine, Maria T.; Humphreys, W. Griffith; Lan, Shih-Jung; Ewing, William R.; Carlson, Kenneth E.; Kowala, Mark C.; Zahler, Robert; Macor, John E.  
CORPORATE SOURCE: Discovery Chemistry and Metabolic and Cardiovascular Drug Discovery, Bristol-Myers Squibb Pharmaceutical Research Institute, Princeton, NJ, 08543-5400, USA  
SOURCE: Journal of Medicinal Chemistry (2005), 48(1), 171-179  
CODEN: JMCMAJ; ISSN: 0022-2623  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
GI

L5 ANSWER 9 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB In a previous report it was demonstrated that merging together key structural elements present in an AT1 receptor antagonist (irbesartan) with key structural elements in a biphenylsulfonamide ETA receptor antagonist followed by addnl. optimization provided a product which is a dual-action receptor antagonist (DARA), which potentially blocked both AT1 and ETA receptors. Described herein are our efforts directed toward improving both the pharmacokinetic profile as well as the AT1 and ETA receptor potency of 4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide (I). Efforts centered on modifying the 2'-side chain of I and examining the (isoxazolyl)sulfonamide moiety in I. This effort resulted in the discovery of 4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-[(ethoxymethyl)[1,1'-biphenyl]-2-sulfonamide (II) as a highly potent second-generation DARA. This compound also showed substantially improved pharmacokinetic properties compared to I. In rats, DARA II reduced blood pressure elevations caused by i.v. infusion of Ang II or big ET-1 to a greater extent and with longer duration than DARA I or AT1 or ETA receptor antagonists alone. II clearly demonstrated superiority over irbesartan (an AT1 receptor antagonist) in the normal SHR model of hypertension in a dose-dependent manner, demonstrating the synergy of AT1 and ETA receptor blockade in a single mol. The crystal and mol. structures of II were reported.  
IT 138402-11-6, Irbesartan  
RL: PAC (Pharmacological activity); BIOL (Biological study)  
(preparation of irbesartan-BMS 193884 combination drug and study of its activity as dual angiotensin II and endothelin A receptor antagonist and its pharmacokinetics)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-[(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)

L5 ANSWER 10 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:696369 CAPLUS  
DOCUMENT NUMBER: 141:225515  
TITLE: Synthesis of 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-1,3-diazaspiro[4.4]non-ene-4-one  
INVENTOR(S): Wissevich, Gennady; Rukhman, Igor; Pertsikov, Boris; Kaftanov, Julia; Dolitzky, Ben-zion  
PATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals Usa, Inc.  
SOURCE: PCT Int. Appl., 27 pp.  
CODEN: PIXOXD  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004072064	A1	20040826	WO 2004-US3604	20040205
W: AE, AE, AG, AL, AL, AM, AM, AT, AU, AZ, BZ, BG, BG, BR, BR, BY, BY, BZ, CA, CH, CH, CO, CO, CR, CR, CU, CU, CZ, CZ, DE, DE, DK, DK, DM, DM, EC, EC, EE, EE, EG, ES, ES, FI, FI, GB, GB, GE, GE, GH, GH, GR, GR, HU, HU, ID, ID, IL, IL, IN, IS, JP, JP, KE, KE, KG, KG, KP, KP, KR, KR, KZ, KZ, LC, LX, LR, LS, LS, LT, LU, LV, MA, MD, MD, MG, MK, MN, MW, MX, MX, MZ, MZ, NA, NI				
RV: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CN, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2004242894	A1	20041202	US 2004-773414	20040205
PRIORITY APPLN. INFO.:			US 2003-445218P	P 20030205
			US 2003-465905P	P 20030428
OTHER SOURCE(S):			CASREACT 141:225515	
GI				



AB Provided are 5 methods of making 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-1,3-diazaspiro[4.4]non-1-ene-4-one (I), e.g. comprising the steps of: (a) reacting 1-(N'-pentanoylamino)cyclopentanecarboxylic acid amide with 5-(4'-bromomethylbiphenyl-2-yl)-1-trityl-1H-

L5 ANSWER 10 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

tetrazole in the presence of an inorg. base, a solvent and a phase transfer catalyst; (b) cooling the mixt.; (c) adding water to the mixt. whereby two phases are obtained; (d) sepg. the two phases obtained; and (e) recovering the compd. I. The compd. I can be converted to irbesartan which is a known angiotensin II receptor antagonist (blocker).  
IT 745814-09-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(methods for preparation of 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-1,3-diazaspiro[4.4]non-ene-4-one)  
RN 745814-09-9 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-(1,1-dimethylethyl)-3-[[2'-(1-(triphenylmethyl)-1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)







L5 ANSWER 11 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:633925 CAPLUS  
DOCUMENT NUMBER: 141:157121  
TITLE: Synthesis of irbesartan  
INVENTOR(S): Nisnevich, Gennady; Rukhman, Igor; Pertsikov, Boris;  
Kaftanov, Julius; Dolitzky, Ben-zion  
PATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva  
Pharmaceuticals Usa, Inc.  
SOURCE: PCT Int. Appl., 21 pp.  
CODEN: PFXK02  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004065383	A2	20040805	WO 2004-US1135	20040116
WO 2004065383	A3	20041216		
V:	AZ, AE, AG, AL, AM, AN, AT, AU, AV, AZ, BA, BB, BG, BR, BS, BY, BZ, CA, CH, CN, CO, CR, CU, CV, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, FO, GB, GD, GE, GF, GH, GI, GR, GT, GU, HK, HN, HU, ID, IL, IN, IS, JP, KE, KG, KH, KI, KP, KR, KZ, LK, LG, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MU, MX, MY, NZ	US 20040930	US 2004-759906	20040116
EP 1509517	A2	20050302	EP 2004-702955	20040116
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK	US 2003-440997P	P 20030116	
PRIORITY APPLN. INFO.:			WO 2004-US1135	W 20040116

OTHER SOURCE(S): CASREACT 141:157121  
AB Provided are a method of making irbesartan via a Suzuki coupling reaction and a novel intermediate, 2-butyl-3-[(4'-bromobenzyl)-1,3-diazaspiro[4.4]non-1-en-4-one, for such process. The novel process includes the step of reacting such intermediate with a protected tetrazolylphenylboronic acid.  
IT 138402-11-6P, Irbesartan  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
RN (Synthesis of irbesartan)  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)




L5 ANSWER 12 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
has renoprotective effect in diabetic nephropathy)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 186 THERE ARE 186 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 12 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:594584 CAPLUS  
DOCUMENT NUMBER: 142:32318  
TITLE: AT1 receptor antagonists: pharmacology  
AUTHOR(S): de Gasparo, M.  
CORPORATE SOURCE: MG Consulting Co, Rossemaison, 2842, Switz.  
SOURCE: Handbook of Experimental Pharmacology (2004), 163/II(Angiotensin, Volume 2), 417-451  
CODEN: HEPH02; ISSN: 0171-2004  
PUBLISHER: Springer-Verlag  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: English  
AB A review. Innovative chemical modifications of the first nonpeptide indazole antagonist of Ang II led to the synthesis of various new orally active agents with increased potency and improved bioavailability (131-801). They block specifically and selectively the angiotensin AT1 receptor without intrinsic agonist properties. The angiotensin receptor blockers (ARB) can be classified as surmountable, (losartan, eprosartan, telmisartan), insurmountable (candesartan) or mixed (valsartan, irbesartan, olmesartan) antagonists depending on their degree of tight binding and their dissociation rate. Candesartan and olmesartan are administered as prodrug converted to the active compound upon absorption. The ARBs are excreted essentially in the bile and mainly unchanged. If biotransformed, it involves oxidative reaction and conjugation. The metabolism of irbesartan, losartan, and candesartan requires cytochrome P 450 enzymes. There is no accumulation by repeated doses. Plasma concns. are little influenced by mild-to-moderate renal impairment but caution may be required in patients with hepatic insufficiency due to the biliary mechanism of excretion. Losartan is unique by its uricosuric property. In general, the ARBs do not interfere with other drugs in a clin. significant way, but caution should be taken if prescribed with potassium-sparing agents or supplements, especially in elderly patients with reduced renal function. The ARBs are generally well tolerated with an incidence of adverse effects or withdrawals similar to the placebo. First-dose hypotension is uncommon and there is no rebound hypertension after withdrawal. Angio-edema is rare. The ARBs are contra-indicated during pregnancy. The efficacy of the ARBs in hypertension is well documented in various population and age groups and better tolerated than other antihypertensive agents for similar efficacy. The first properly powered trial in an hypertensive population, LIFE with losartan, has demonstrated a beneficial effect on the primary composite endpoint including cardiovascular death, myocardial infarction and stroke, even more impressive in a diabetic subgroup. Based on the result of Val-HeFT, valsartan is approved in heart failure patients intolerant to ACE inhibitor. The renoprotective effect of the ARBs was demonstrated in diabetic nephropathy with irbesartan, losartan and valsartan. Various clin. studies suggest a beneficial effect of the ARBs beyond the blood pressure fall. Several large trials are in progress to establish the efficacy of the ARBs in patients with LV dysfunction following recent myocardial infarction. As blockade of the AT1 receptor is accompanied by increased plasma Ang II, the potential of the stimulation of the unblocked AT2 receptor is discussed. Possible further indications for ARB are also briefly reported.  
IT 138402-11-6, Irbesartan  
RL: ADV (Adverse effect, including toxicity); DMA (Drug mechanism of action); PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(mixed ARB irbesartan requires cytochrome P 450 enzymes for metabolism and

L5 ANSWER 13 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:475010 CAPLUS  
DOCUMENT NUMBER: 141:405861  
TITLE: Effect of irbesartan on angiotensin II-induced hypertrophy of human proximal tubular cells  
AUTHOR(S): Liu, Bi-cheng; Sun, Jing; Chen, Qi; Luo, Dong-dong; Ma, Kun-ling; Ruan, Xiong-zhong  
CORPORATE SOURCE: Institute of Nephrology, Zhongda Hospital, Southeast University School of Medicine, Nanjing, 210009, Peop. Rep. China  
SOURCE: Chinese Medical Journal (Beijing, China, English Edition) (2004), 117(4), 547-551  
CODEN: CMJ005; ISSN: 0366-6999  
PUBLISHER: Chinese Medical Association  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Background: Intrarenal activation of the renin angiotensin system (RAS) plays an important role in mediating renal fibrosis. Both angiotensin converting enzyme inhibitors (ACEIs) and angiotensin II (Ang II) receptor antagonists have been shown to exert a protective role against diabetic and non-diabetic nephropathy. However, the exact mechanism of how blocking local RAS prevents renal fibrosis is unclear. The present study was to investigate the influence of a new Ang II receptor antagonist, irbesartan (Irb), on Ang II-induced hypertrophy in human proximal tubular cell line (HK-2). Methods: The cell line, HK-2, was grown in Dulbeccos's Modified Eagle's Medium containing 10% heat-inactivated fetal calf serum. After rested in serum-free medium for 24 h, the effects of Irb on Ang II [10<sup>-7</sup> mol/L]-induced [3H]-leucine incorporation, total protein content (measured by the Coomassie brilliant blue G250 method), and change in cell size (determined by SEM) were observed. The influence of Irb on the cell cycle was analyzed by fluorescence activated cell sorter (FACS) flow cytometry. Results: Ang II induced cell hypertrophy in a time and dose dependent manner. Stimulation of cells with Ang II for 48 h resulted in an increase in [3H]-leucine incorporation [0 h (5584 ± 1016) cpm/105 cells vs 48 h: (10741 ± 802) cpm/105 cells, P < 0.05], which was significantly attenuated by treatment with Irb. Ang II significantly increased the total protein content in HK-2 cells [control: (0.169 ± 0.011) μg/105 cells vs Ang II group: (0.202 ± 0.010) μg/105 cells, P < 0.05], which was also markedly inhibited by cotreatment with Irb (P < 0.01). SEM showed that Ang II induced an increase in average phys. cell size, which was significantly inhibited by Irb [control: (11.92 ± 1.62) μm; Ang II group: (20.63 ± 3.83) μm; Ang II + Irb group: (13.59 ± 3.15) μm; P < 0.01 vs control, resp.]. Furthermore, flow cytometry revealed that Ang II arrested cells in the G0-G1 phase, which was significantly reversed by treatment with Irb [G0-G1 cells in Ang II group: (76.09 ± 0.92)%, in Ang II + Irb group: (67.00 ± 2.52)%, P < 0.05]. Conclusion: Irb can inhibit Ang II-induced hypertrophy in HK-2 cells.  
IT 138402-11-6, Irbesartan  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(angiotensin II receptor antagonist irbesartan inhibited Ang II-induced hypertrophy of human proximal tubular HK-2 cell line)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


ACCESSION NUMBER: 2004:451474 CAPLUS  
DOCUMENT NUMBER: 141:1258  
TITLE: Nitrosated compounds in methods of treating vascular diseases characterized by nitric oxide insufficiency  
INVENTOR(S): Loscalzo, Joseph; Vita, Joseph A.; Loberg, Michael D.; Worcel, Manuel  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 23 pp., Cont.-in-part of U.S. Ser. No. 679,257.  
CODEN: USXKXO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 5  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004105850	A1	20040603	US 2003-692724	20031027
US 6635273	B1	20031021	US 2000-697317	20001027
US 2004071766	A1	20040415	US 2003-679257	20031007
PRIORITY APPLN. INFO.:			US 1999-162230P	P 19991029
			US 2000-179020P	P 20000131
			US 2000-697317	A1 20001027
			US 2003-679257	A2 20031007

OTHER SOURCE(S): MARPAT 141:1258  
AB The invention provides methods of treating and/or preventing vascular diseases characterized by nitric oxide insufficiency by administering a therapeutically effective amount of at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated cholesterol reducer, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and optionally at least one compound used to treat cardiovascular diseases and/or at least one antioxidant, or a pharmaceutically acceptable salt thereof, and/or at least one compound that donates, transfers or releases nitric oxide, elevates endogenous levels of endothelium-derived relaxing factor, stimulates endogenous synthesis of nitric oxide or is a substrate for nitric oxide synthase. The antioxidant may preferably be a hydralazine compound or a pharmaceutically acceptable salt thereof. The compound that donates, transfers or releases nitric oxide, elevates endogenous levels of endothelium-derived relaxing factor, stimulates endogenous synthesis of nitric oxide or is a substrate for nitric oxide synthase may preferably be isosorbide dinitrate and/or isosorbide mononitrate. The vascular diseases characterized by nitric oxide insufficiency include a cardiovascular disease and a disease resulting from oxidative stress. Nitric oxide action was shown to be impaired in the microvasculature of black hypertensive patients to a greater extent than in white hypertensive patients.  
IT 138402-11-6D, Irbesartan, nitrosated compds.  
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(nitrosated compds. in methods of treating vascular diseases characterized by nitric oxide insufficiency)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



ACCESSION NUMBER: 2004:404854 CAPLUS  
DOCUMENT NUMBER: 140:417642  
TITLE: Diverse effects of long-term treatment with imidapril and irbesartan on cell growth signal, apoptosis and collagen type I expression in the left ventricle of spontaneously hypertensive rats  
AUTHOR(S): Wang, Jin-ming; Wang, Ying; Zhu, Zhong-sheng; Zhang, Mei-chun; Zou, Yi; Li, Jian-jun; Li, Ming-jiang; Jiang, Xue-jun; Li, Xiao-Yan  
CORPORATE SOURCE: Renmin Hospital, Department of Cardiology, Wuhan University School of Medicine, Wuhan, 430060, Peop. Rep. China  
SOURCE: Life Sciences (2004), 75(4), 407-420  
CODEN: LIFSAK; ISSN: 0024-3205  
PUBLISHER: Elsevier  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB To compare diverse effects of angiotensin II type 1 receptor antagonists (irbesartan) and angiotensin converting enzyme inhibitors (imidapril) on left ventricular remodeling in spontaneously hypertensive rats (SHR). Thirty male SHR were randomly divided into three groups: SHR-IR (treated with irbesartan, 50 mg/kg), SHR-IM (imidapril, 3 mg/kg), SHR-C (placebo). Ten male Wistar Kyoto rats (WKY) treated with placebo acted as the control. All treatments were administered once daily from 14 to 27 wk of age. Imidapril and irbesartan have the similar inhibitor effects on blood pressure and left ventricular mass indexes in SHR. Despite both drugs suppressed ERK-1 protein expression, decreased cardiomyocytes apoptosis index, blocked collagen type I deposition, reduced TGF-β1 gene expression in SHR, imidapril elicits a stronger inhibitory effect. Irbesartan had little effect on MKP-1 protein expression, but imidapril decreased it significantly. As a result, the ERK-1/MKP-1 ratio in SHR-IR was significantly greater than that in SHR-IM (P < 0.05). These results suggest that the balance between MKP-1 and ERKs in myocardial tissue is important for cardiac cell proliferation and growth. They also indicate that the similar efficacy of antihypertensive treatment in reducing blood pressure does not predict the similar capacity to control the individual facet of left ventricular remodeling. Irbesartan is more effective in regressing the homeostasis between ERK-1 and MKP-1, however imidapril is superior in suppressing apoptosis and collagen synthesis in cardiac tissue.  
IT 138402-11-6, Irbesartan  
RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(diverse effects of long-term treatment with imidapril and irbesartan on cell growth signal, apoptosis and collagen I in left ventricle in hypertension)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 15 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 16 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:392331 CAPLUS  
DOCUMENT NUMBER: 140:406798  
TITLE: Preparation of benzoxepinopyridines as HMG-CoA  
reductase inhibitors  
INVENTOR(S): Robl, Jeffrey A.; Chen, Bang-chi; Sun, Chong-qing  
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
SOURCE: U.S. Pat. Appl. Publ., 44 pp., Cont.-in-part of U.S.  
Ser. No. 875,155, abandoned.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 3  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004092573	A1	20040513	US 2003-602752	20030624
US 6812345	B2	20041102		
US 2002013334	A1	20020131	US 2001-875155	20010606
PRIORITY APPLN. INFO.:			US 2000-211595P	P 20000615
			US 2001-875155	B2 20010606

OTHER SOURCE(S): MARPAT 140:406798  
GI


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I [X = O, S, SO, SO<sub>2</sub>, NR<sub>7</sub>; 2 = HOCHCH<sub>2</sub>CH(OH)CH<sub>2</sub>CO<sub>2</sub>R<sub>3</sub>,  
4-hydroxy-2-oxopyran-6-yl, etc.; n = 0, 1; R<sub>1</sub>, R<sub>2</sub> = alkyl, arylalkyl,  
cycloalkyl, alkenyl, cycloalkenyl, aryl, heteroaryl, cycloheteroalkyl; R<sub>3</sub>  
= H, alkyl, metal ion; R<sub>4</sub> = H, halo, CF<sub>3</sub>, etc.; R<sub>7</sub> = H, alkyl, aryl,  
alkenyl, acryl, alkoxycarbonyl, etc.; R<sub>9</sub>, R<sub>10</sub> = H, alkyl], were prepared as  
HMG CoA reductase inhibitors active in inhibiting cholesterol  
biosynthesis, modulating blood serum lipids such as lowering LDL  
cholesterol and/or increasing HDL cholesterol, and treating  
hyperlipidemia, hypercholesterolemia, hypertriglyceridemia and  
atherosclerosis (no data). A multistep synthesis of II is  
reported.

IT 138402-11-6, Irbesartan  
RI: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
(Biological study); USES (Uses)  
(coadministered agents; preparation of benzoxepinopyridines as HMG-CoA  
reductase inhibitors for treatment of hyperlipidemia,  
hypercholesterolemia, hypertriglyceridemia, atherosclerosis, and other  
disorders)

RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-  
biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)

L5 ANSWER 16 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 17 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:292572 CAPLUS  
DOCUMENT NUMBER: 141:16887  
TITLE: Novel dual action AT<sub>1</sub> and ETA receptor antagonists  
reduce blood pressure in experimental hypertension  
Kowala, Mark C.; Murugesan, Natesan; Taliev, John;  
Carlson, Kenneth; Monshizadeh, Hossein; Ryan, Carol;  
Gu, Zhengxiang; Kane, Bridgette; Fadnis, Leena; Baska,  
Rose Ann; Beyer, Sophie; Arthur, Susan; Dickinson,  
Kenneth; Zhang, Donglu; Perrone, Mark; Ferrer, Pam;  
Giancarli, Mary; Baumann, Jergen; Bird, Eileen;  
Panchal, Balkrushna; Yang, Yifan; Trippodo, Nick;  
Barrish, Joel; Macor, John E.  
Departments of Metabolic and Cardiovascular Drug  
Discovery, Bristol-Myers Squibb Pharmaceutical  
Research Institute, Princeton, NJ, USA  
SOURCE: Journal of Pharmacology and Experimental Therapeutics  
(2004), 309(1), 275-284  
CODEN: JPETAB; ISSN: 0022-3565  
PUBLISHER: American Society for Pharmacology and Experimental  
Therapeutics  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Angiotensin II and endothelin-1 activate their resp. AT<sub>1</sub> and ETA receptors  
on vascular smooth muscle cells, producing vasoconstriction, and both  
peptides are implicated in the pathogenesis of essential hypertension.  
Angiotensin II potentiates the production of endothelin, and conversely  
endothelin augments the synthesis of angiotensin II. Both AT<sub>1</sub>  
and ETA receptor antagonists lower blood pressure in hypertensive  
patients; thus, a combination AT<sub>1</sub>/ETA receptor antagonist may have greater  
efficacy and broader utility compared with each drug alone. By rational  
drug design a biphenyl ETA receptor blocker was modified to acquire AT<sub>1</sub>  
receptor antagonism. These compds. (C and D) decreased  
Sar-Ile-Angiotensin II binding to AT<sub>1</sub> receptors and endothelin-1 binding  
to ETA receptors, and compound C inhibited angiotensin II- and  
endothelin-1-mediated Ca<sup>2+</sup> transients. In rats compds. C and D reduced  
blood pressure elevations caused by i.v. infusion of angiotensin II or big  
endothelin-1. Compound C decreased blood pressure in Na<sup>+</sup>-depleted  
spontaneously hypertensive rats and in rats with mineralocorticoid  
hypertension. Compound D was more efficacious than AT<sub>1</sub> receptor antagonists  
at reducing blood pressure in spontaneously hypertensive rats, and its  
superiority was likely due to its partial blockade of ETA receptors.  
Therefore compds. C and D are novel agents for treating a broad spectrum  
of patients with essential hypertension and other cardiovascular diseases.

IT 138402-11-6, Irbesartan  
RI: PAC (Pharmacological activity); BIOL (Biological study)  
(novel dual action AT<sub>1</sub> and ETA receptor antagonists reduce blood  
pressure in exptl. hypertension)

RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-  
biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2004:60496 CAPLUS  
DOCUMENT NUMBER: 140:111420  
TITLE: Synthesis of irbesartan  
INVENTOR(S): Nisnevich, Gennady; Rukhman, Igor; Pertsikov, Boris; Kaftanov, Julia; Dolitzky, Ben-Zion; Shapiro, Eugeny; Yahalomi, Bonit  
PATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals USA, Inc.  
SOURCE: PCT Int. Appl., 13 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007482	A2	20040122	WO 2003-US22479	20030716
WO 2004007482	A3	20040527		
V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RV: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2492779	AA	20040122	CA 2003-2492779	20030716
EP 1546135	A2	20050629	EP 2003-764805	20030716
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, IL, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
US 2005176794	A1	20050811	US 2003-621623	20030716
PRIORITY APPLN. INFO.: US 2002-396424P P 20020716				
US 2002-402490P P 20020809				
WO 2003-US22479 W 20030716				

OTHER SOURCE(S): CASREACT 140:111420  
AB Irbesartan is prepared by reaction of 2-butyl-1,3-diaza-spiro[4.4]non-1-ene (I) with 5-(4-bromomethylbiphenyl-2-yl)-1H-tetrazole (II) in the presence of a phase transfer catalyst. Thus, reaction of I with II in toluene in the presence of Bu4NHSO4 at 90° for 1.5 h gave, after deprotection, 84.3% irbesartan. Also provided is irbesartan having a fine particle size.  
IT 138402-11-6P, Irbesartan  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




ACCESSION NUMBER: 2003:696766 CAPLUS  
DOCUMENT NUMBER: 139:207790  
TITLE: Antihypertensive composition and method using an antihypertensive agent-antiinflammatory agent combination  
INVENTOR(S): Hamet, Pavel; Tremblay, Johanne  
PATENT ASSIGNEE(S): Corporation du Centre de Recherche du Chum, Can.  
SOURCE: PCT Int. Appl., 46 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003072136	A1	20030904	WO 2003-CA266	20030226
V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RV: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.: US 2002-359331P P 20020226				
AB A method and compns. are provided for attenuating and/or preventing blood pressure increment caused by a stressful stimuli. The method consists in administering to a mammal a composition including an effective amount of at least one antihypertensive compound in combination with an effective amount of at least one antiinflammatory agent.				
IT 138402-11-6, Irbesartan				
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
(antihypertensive agent-antiinflammatory agent combination for stress-caused hypertension)				
RN 138402-11-6 CAPLUS				
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)				



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 20 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:587858 CAPLUS  
 DOCUMENT NUMBER: 137:288496  
 TITLE: Discovery of N-isoxazolyl biphenylsulfonamides as potent dual angiotensin II and endothelin A receptor antagonists  
 AUTHOR(S): Murugesan, Natesan; Tellez, John E.; Gu, Zhengxiang; Kunst, Bridgette L.; Fadnis, Leena; Cornelius, Lyndon A.; Baska, Rose Ann F.; Yang, Yifan; Beyer, Sophie M.; Monshizadegan, Hossain; Dickinson, Kenneth E.; Panchal, Balkrushna; Valentine, Maria T.; Chong, Saeho; Morrison, Richard A.; Carlson, Kenneth E.; Powell, James R.; Moreland, Suzanne; Barish, Joel C.; Kowala, Mark C.; Macor, John E.  
 CORPORATE SOURCE: Discovery Chemistry and Metabolic and Cardiovascular Drug Discovery, Bristol-Myers Squibb Pharmaceutical Research Institute, Princeton, NJ, 08543-5400, USA  
 SOURCE: Journal of Medicinal Chemistry (2002), 45(18), 3829-3835  
 CODEN: JMCHAR; ISSN: 0022-2623  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 137:288496  
 AB The ETA receptor antagonist N-(3,4-dimethyl-5-isoxazolyl)-4'-[(2-oxazolyl)-[1,1'-biphenyl]-2-sulfonamide (BMS-193884) shares the same biphenyl core as a large number of AT1 receptor antagonists, including irbesartan. Thus, it was hypothesized that merging the structural elements of BMS-193884 with those of the biphenyl AT1 antagonists (e.g., irbesartan) would yield a compound with dual activity for both receptors. This strategy led to the design, synthesis, and discovery of 4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-[1,1'-biphenyl]-2-sulfonamide (BMS-248360) as a potent and orally active dual antagonist of both AT1 and ETA receptors. Compound BMS-248360 represents a new approach to treating hypertension.  
 IT 138402-11-6, Irbesartan  
 RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study)  
 (discovery of N-isoxazolyl biphenylsulfonamides as potent dual angiotensin II and endothelin A receptor antagonists)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 22 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:540258 CAPLUS  
 DOCUMENT NUMBER: 137:109267  
 TITLE: Preparation of benzoxepinopyridines as HMG-CoA reductase inhibitors  
 INVENTOR(S): Robl, Jeffrey A.; Chen, Bang-chi; Sun, Chong-qing  
 PATENT ASSIGNER(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 42 pp., Cont.-in-part of U.S. Ser. No. 875,155.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002094977	A1	20020718	US 2001-7407	20011204
US 6627636	B2	20030930		
US 2002013334	A1	20020131	US 2001-875155	20010606
PRIORITY APPLN. INFO.:			US 2000-211595P	P 20000615
			US 2001-875155	A2 20010606

OTHER SOURCE(S): MARPAT 137:109267  
 GI


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I [X = O, S, SO, SO<sub>2</sub>, NR<sub>7</sub>; Z = HOCH<sub>2</sub>CH<sub>2</sub>CH(OH)CH<sub>2</sub>CO<sub>2</sub>R<sub>3</sub>, 4-hydroxy-2-oxopyran-6-yl, etc.; n = 0, 1; R<sub>1</sub>, R<sub>2</sub> = alkyl, arylalkyl, cycloalkyl, alkenyl, cycloalkenyl, aryl, heteroaryl, cycloheteroalkyl; R<sub>3</sub> = H, alkyl, metal ion; R<sub>4</sub> = H, halo, CF<sub>3</sub>, etc.; R<sub>7</sub> = H, alkyl, aryl, alkanoyl, acryl, alkoxycarbonyl, etc.; R<sub>9</sub>, R<sub>10</sub> = H, alkyl], were prepared as HMG CoA reductase inhibitors active in inhibiting cholesterol biosynthesis, modulating blood serum lipids such as lowering LDL cholesterol and/or increasing HDL cholesterol, and treating hyperlipidemia, hypercholesterolemia, hypertriglyceridemia and atherosclerosis (no data). A multistep synthesis of II is reported.  
 IT 138402-11-6, Irbesartan  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (coadministered agents; preparation of benzoxepinopyridines as HMG-CoA reductase inhibitors for treatment of hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, atherosclerosis, and other disorders)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)

L5 ANSWER 21 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:569152 CAPLUS  
 DOCUMENT NUMBER: 138:104378  
 TITLE: Forefront of diabetes and renal diseases  
 AUTHOR(S): Kamiya, Yoshinobu; Kimura, Genjiro  
 CORPORATE SOURCE: Graduate School of Medicine, Nagoya City University, Japan  
 SOURCE: Bunshi Shin Kekkanyo (2002), 3(2), 209-215  
 CODEN: BSKUAB; ISSN: 1345-2355  
 PUBLISHER: Sentan Igakusha  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: Japanese  
 AB A review on role of angiotensin II in diabetic nephropathy. The topics discussed are (1) renal renin-angiotensin system in diabetic nephropathy; (2) glucose-induced angiotensin II in mesangial cells; (3) glucose and angiotensin II in stimulating extracellular matrix protein synthesis through induction of transforming growth factor-β expression; (4) association of genetic polymorphisms of proteins in renin-angiotensin system with diabetic nephropathy; and (5) effects of angiotensin II receptor blockers losartan and irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes.  
 IT 138402-11-6, Irbesartan  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (angiotensin II and its receptor in diabetic nephropathy and role of angiotensin receptor antagonists)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




L5 ANSWER 22 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



L5 ANSWER 23 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:392237 CAPLUS  
 DOCUMENT NUMBER: 136:401651  
 TITLE: Preparation of fused pyridine derivatives as HMG-CoA reductase inhibitors  
 INVENTOR(S): Robl, Jeffrey A.; Chen, Bang-Chi; Sun, Chong-Qing  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 46 pp., Cont.-in-part of U.S. Ser. No. 875,218.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002061901	A1	20020523	US 2001-8154	20011204
US 6620821	B2	20030916		
US 2002028826	A1	20020307	US 2001-875218	20010606
US 2004024216	A1	20040205	US 2003-602753	20030624
			US 2000-211594P	P 20000615
			US 2001-875218	A2 20010606
			US 2001-8154	A3 20011204

PRIORITY APPLN. INFO.:  
 OTHER SOURCE(S): MARPAT 136:401651  
 GI



AB The title compds. I and their pharmaceutically acceptable salts, esters, prodrug esters, and stereoisomers are claimed [wherein: Z = CH(OH)CH2CR7(OH)CH2CO2R3 or corresponding pyranone lactone deriva., n = 0, 1; m = 0, 1, 2, 3, or 4; y = 0, 1, 2, 3 or 4, provided that at least one of x and y is other than 0; and optionally one or more carbons of (CH2)x and/or (CH2)y together with addnl. carbons form a 3 to 7 membered spirocyclic ring; R1, R2 = alkyl, arylalkyl, cycloalkyl, alkenyl, cycloalkenyl, aryl, heteroaryl, cycloheteroalkyl; R3 = H or lower alkyl; R4 = H, halo, CF3, OR, alkyl, alkoxy, CO2R, (un)substituted NH2, cyano, (un)substituted CONH2, etc.; R7 = H, alkyl]. The compds. are HMG-CoA reductase inhibitors, and are active in inhibiting cholesterol biosynthesis and modulating blood serum lipids, for example, lowering LDL cholesterol and/or increasing HDL cholesterol (no data). I are thus useful in treating hyperlipidemia and dyslipidemia, in hormone replacement therapy, and in treating hypercholesterolemia, hypertriglyceridemia and atherosclerosis, as well as Alzheimer's disease and osteoporosis. Prepsns.

L5 ANSWER 24 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:315368 CAPLUS  
 DOCUMENT NUMBER: 136:330577  
 TITLE: Tissue perfusion enhancement by co-administration of a drug that increases cGMP synthesis and an agent that inhibits cGMP degradation  
 INVENTOR(S): Mueller, Thomas H.  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 3 pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002048599	A1	20020425	US 2001-981335	20011016
WO 2002034248	A2	20020502	WO 2001-US42742	20011016
WO 2002034248	A3	20030403		

PRIORITY APPLN. INFO.:  
 AB A method for increasing tissue perfusion with blood by the co-administration of an agent that increases cGMP synthesis and an agent that inhibits cGMP degradation in the cells of the blood vessel walls


or in blood cells. The method comprises, for example, the co-administration of a statin and dipyradimole, especially a timed-release formulation of dipyradimole. Atorvastatin at 5-80 or fluvastatin at 10-40 mg/day can be used.

IT 138402-11-6, Irbesartan  
 RI: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (tissue perfusion enhancement by co-administration of drugs that increase cGMP synthesis)

RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




L5 ANSWER 23 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 of several compds. are described. For instance, a multistep synthesis of fused pyridine deriv. II is reported. Compds. I may be used in a manner similar to atorvastatin, pravastatin, simvastatin, etc. Combinations of compds. I with various other drugs are claimed, the latter being specified as certain pharmacol. classes, as inhibitors of specific enzymes, as (ant)agonists of specific receptors, and as numerous named drugs.  
 IT 138402-11-6, Irbesartan  
 RI: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (therapeutic compns. also containing; preparation of fused pyridine deriva. as HMG-CoA reductase inhibitors)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 25 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2002:203409 CAPLUS  
 DOCUMENT NUMBER: 136:353534  
 TITLE: Lack of impairment of nitric oxide-mediated responses in a rat model of high-renin hypertension  
 AUTHOR(S): Artigues-Varin, C.; Richard, V.; Renet, S.; Henry, J. P.; Thuilliez, C.  
 CORPORATE SOURCE: Department of Pharmacology, INSERM EMI 9920, IFRMP no 23, Rouen University Medical School, Rouen, 76183, Fr.  
 SOURCE: Clinical and Experimental Pharmacology and Physiology (2002), 29(1/2), 26-31  
 CODEN: CEXPB9; ISSN: 0305-1870  
 PUBLISHER: Blackwell Publishing Asia  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English


AB 1. Angiotensin (Ang) II triggers the expression of a prooxidant phenotype in the vascular wall, suggesting that activation of the renin-angiotensin system (RAS) causes endothelial dysfunction in various pathol. situations, such as hypertension. However, this hypothesis has been mostly tested in a setting of exogenous administration of AngII. 2. We tested the hypothesis of a role for endogenous activation of the RAS leading to oxidant stress and endothelial dysfunction in a high-renin model of hypertension (i.e. two-kidney, one-clip hypertension) in rats. One month after clipping or sham surgery, aorta were isolated from untreated rats or rats treated by the angiotensin AT1 receptor antagonist irbesartan (10 mg/kg per day). Mesenteric artery segments were also isolated from normotensive or hypertensive rats. 3. Hypertension reduced the relaxations to acetylcholine but did not affect the ratio of contractions to phenylephrine in the presence compared with the absence of a nitric oxide (NO) synthase inhibitor, used as an index of basal release of NO. 4. The free radical scavenger tempol reduced the contractions to phenylephrine in the absence, but not in the presence, of an inhibitor of NO synthesis. This index of free radical-mediated degradation of NO was not affected by hypertension. In parallel, hypertension did not affect the expression of p22phox, a component of the free radical generating enzyme reduced NADPH oxidase. 5. Chronic treatment with the AT1 receptor antagonist decreased blood pressure, moderately improved the response to acetylcholine, but did not affect basal NO release in hypertensive rats, although it did increase basal NO release in normotensive rats. 6. Thus, this model of hypertension is characterized by an impaired stimulated NO release but not of basal NO release in isolated arteries. Furthermore, there was no functional evidence of an increased oxidative stress-mediated impairment of NO release. This is not in favor of a direct link between activation of the RAS and development of endothelial dysfunction in exptl. hypertension.

IT 138402-11-6, Irbesartan  
 RI: BSU (Biological study, unclassified); BIOL (Biological study)  
 (lack of impairment of nitric oxide-mediated responses in a rat model of high-renin hypertension)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2001:426040 CAPLUS  
DOCUMENT NUMBER: 135:283303  
TITLE: Synthesis and study of a cyclic angiotensin II antagonist analogue reveals the role of  $\pi$ - $\pi$  interactions in the C-terminal aromatic residue for agonist activity and its structure resemblance with AT1 non-peptide antagonists  
AUTHOR(S): Polevaya, L.; Mavroussakos, T.; Zomboulakis, P.; Grdadolnik, S. G.; Roumelioti, P.; Giatas, N.; Mutula, I.; Keivish, T.; Vlahakos, D. V.; Iliodromitis, E. K.; Kremastinos, D. Th.; Matsoukas, J.  
CORPORATE SOURCE: Laboratory of Peptide Chemistry, Latvian Institute of Organic Synthesis, Riga, LV-1006, Latvia  
SOURCE: Bioorganic & Medicinal Chemistry (2001), 9(6), 1639-1647  
CODEN: BMECEP; ISSN: 0968-0896  
PUBLISHER: Elsevier Science Ltd.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The novel amide linked Angiotensin II (ANG II) cyclic analog cyclo(3,5)-[Sar1-Lys3-Glu5-Ile8] ANG II (18) has been designed, synthesized and bioassayed in anesthetized rabbits. The constrained cyclic analog with a lactam amide bridge linking a Lys-Glu pair at positions 3 and 5 and possessing Ile at position 8, was synthesized by solution procedure using the maximum protection strategy. This analog was found to be inhibitor of Angiotensin II. NMR spectroscopy coupled with computational anal. showed clustering between the side chains of the key amino acids Tyr4-His6-Ile8 similar to that observed with ANG II. The obtained data show that only  $\pi$ - $\pi$  interactions observed in ANG II or its superagonist Sar1 [ANG II] are missing. Therefore, it can be concluded that these interactions are essential for agonist activity. Conformational anal. comparisons between AT1 antagonists losartan, eprosartan and irbesartan with C-terminal segment of cyclic compound 18 revealed structural similarities.  
IT 138402-11-6, Irbesartan  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(synthesis of a cyclic angiotensin II antagonist analog reveals role of  $\pi$ - $\pi$  interactions in C-terminal aromatic residue for agonist activity and structure resemblance with AT1 non-peptide antagonists)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2001:405278 CAPLUS  
DOCUMENT NUMBER: 136:183766  
TITLE: Improvement on synthetic technology of irbesartan  
AUTHOR(S): Shen, Jingshan; Yan, Tiema; Li, Huijun; Li, Jianfeng; Lei, Lijun; Ji, Ruyun  
CORPORATE SOURCE: Department of Medicinal Chemistry, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, 200031, Peop. Rep. China  
SOURCE: Zhongguo Yaowu Huaxue Zazhi (2001), 11(2), 104-106  
CODEN: ZYHZEJ; ISSN: 1005-0108  
PUBLISHER: Zhongguo Yaowu Huaxue Zazhi Bianjibu  
DOCUMENT TYPE: Journal  
LANGUAGE: Chinese  
OTHER SOURCE(S): CASREACT 136:183766  
AB Two methods for synthesizing irbesartan were presented. Irbesartan was prepared from 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one (I) and 4-bromomethyl-2'-cyanobiphenyl by substitution and cyclization with a good yield of 63%. It can also be obtained from I and 2-(4'-bromo-1,1'-biphenyl-2-yl)-2-triphenylmethyltetrazole through substitution and deprotection with overall yield of 85%. The structure was confirmed by 1H-NMR and MS.  
IT 138402-11-6P, Irbesartan  
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(synthesis of irbesartan)  
RN 138402-11-6 CAPLUS  
CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



ACCESSION NUMBER: 2001:323259 CAPLUS  
 DOCUMENT NUMBER: 135:267028  
 TITLE: Renal safety of combined cyclooxygenase 2 (COX-2) inhibitor and angiotensin II receptor blocker administration in mild volume depletion  
 AUTHOR(S): Kistler, Thomas; Ambuhl, Patrice M.  
 CORPORATE SOURCE: Renal Division, University Hospital, Zurich, Switz.  
 SOURCE: Swiss Medical Weekly (2001), 131(13/14), 193-198  
 CODEN: SMWAI; ISSN: 1424-7860  
 PUBLISHER: EMS Swiss Medical Publishers Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Principles: Drugs that either inhibit prostaglandin synthesis or antagonize angiotensin II effects are likely to impair renal function, especially in patients with an activated renin-angiotensin-aldosterone system.


Of the former, non-steroidal anti-inflammatory drugs (NSAIDs) are widely used, and newer agents with cyclooxygenase 2 (COX-2) specific inhibition may have fewer renal side effects compared to non-selective NSAIDs. We therefore investigated whether combination of a COX-2 inhibitor with an angiotensin II subtype 1 (AT1) receptor blocker is safe with regard to preservation of normal renal function in a state of slight volume contraction. Methods: Mild volume depletion was induced by a salt-restricted diet in 5 healthy volunteers who were then given a single dose of 400 mg celecoxib, a COX-2 inhibitor, alone or in combination with 150 mg irbesartan, an AT1 receptor blocker. Glomerular filtration rate (GFR) and effective renal plasma flow (ERPF) were determined by measuring inulin and PAH clearance resp., along with plasma renin activity (PRA) and urinary electrolyte excretion before and over 100 min after drug administration. Results: PRA was high prior to drug administration, indicating slight salt depletion, and dropped by 65% after intake of celecoxib alone (p = 0.008) but only by 25% after combined intake with irbesartan (p = n.s.). GFR was not affected either by celecoxib alone or by combined administration with irbesartan. In contrast, ERPF increased by 28% 80 min after simultaneous drug intake (p = 0.029), but not after celecoxib alone. Renal sodium and potassium excretion did not significantly change under celecoxib alone or in combination with irbesartan. Conclusion: Selective COX-2 inhibition by celecoxib in combination with an AT1 receptor blocker (irbesartan) has no acute adverse effects on renal hemodynamics and renal salt handling in slightly volume-depleted subjects with normal renal function. Moreover, our data obtained in humans appear to confirm the co-regulatory interaction of COX-2 and angiotensin II in the control of renin release, as suggested by animal studies.

17 138402-11-6, Irbesartan

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(renal safety of combined cyclooxygenase 2 (COX-2) inhibitor and angiotensin II receptor blocker administration in mild volume depletion)

RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THIS RE FORMAT

ACCESSION NUMBER: 2001:283949 CAPLUS  
 DOCUMENT NUMBER: 134:311218  
 TITLE: Synthesis and use of heterocyclic sodium/proton exchange inhibitors  
 INVENTOR(S): Ahmad, Saleem; Wu, Shung C.; O'Neil, Steven V.; Ngu, Khehyong; Atwal, Karnail S.  
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
 SOURCE: PCT Int. Appl., 221 pp.  
 CODEN: PIXKX2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001027107	A2	20010419	WO 2000-US27461	20001002
WO 2001027107	A3	20020124		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 6887870	B1	20050503	US 2000-669298	20000925
CA 2388813	AA	20010419	CA 2000-2388813	20001002
EP 1224183	A2	20020724	EP 2000-968723	20001002
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, NO, MK, CY, AL				
BR 2000014725	A	20030617	BR 2000-14725	20001002
JP 2003527331	T2	20030916	JP 2001-530325	20001002
NZ 517668	A	20040924	NZ 2000-517668	20001002
ZA 2002002479	A	20040727	ZA 2002-2479	20020327
NO 2002001717	A	20020610	NO 2002-1717	20020411
US 2005137216	A1	20050623	US 2005-46993	20050131
PRIORITY APPLN. INFO.:			US 1999-158755P	P 19991012
			US 2000-669298	A3 20000925
			WO 2000-US27461	W 20001002

OTHER SOURCE(S): MARPAT 134:311218  
 GI



II

AB Comps. of formula I [wherein: n is 1-5; X is N or CR5, where R5 is H, halo, alkenyl, alkynyl, alkoxy, alkyl, aryl or heteroaryl; Z is a heteroaryl group; R1 is H, alk(en)yl, alk(enyl)ynyl, alkoxy, aryl or

alkyl)3Si, cycloalk(en)yl, (aryl)amino, aryl(alkyl), cycloheteroaryl, etc.; R2, R3 and R4 are any of the groups set out for R1 and optionally substituted with 1 to 5 substituents which may be the same or different and when X is N, R1 is preferably aryl or heteroaryl) are claimed. Several hundred examples are disclosed. Synthesis of II proceeds via cyclopropanation of the cinnamate derived from the olefination between 3,5-dichlorobenzaldehyde and t-butylidenebisphosphonate. The intermediate tert-Bu ester is converted to the corresponding α-chloroketone and reacted with acetyl guanidine to provide II in a total of 5 steps. Comps. I are said to be sodium/proton exchange inhibitors (NHE). Pharmaceutical combinations are claimed using I and certain antihypertensive agents, β-adrenergic agonists, hypolipidemic agents, antidiabetic agents, antiobesity agents, etc. Comps. I are useful as antianginal and cardioprotective agents and provide a method for preventing or treating angina pectoris, cardiac dysfunction, myocardial necrosis, and arrhythmia.


17 138402-11-6, Irbesartan

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(pharmaceuticals also containing; synthesis and use of heterocyclic sodium/proton exchange inhibitors)


RN 138402-11-6 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)






L5 ANSWER 30 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
 ACCESSION NUMBER: 2001:79544 CAPLUS  
 DOCUMENT NUMBER: 134:261085  
 TITLE: ACE inhibitor and AT1 antagonist blockade of deformation-induced gene expression in the rabbit jugular vein through B2 receptor activation  
 AUTHOR(S): Lauth, Manfred; Cattaruzza, Marco; Hecker, Markus  
 CORPORATE SOURCE: Department of Cardiovascular Physiology, University of Göttingen, Göttingen, 37073, Germany  
 SOURCE: Arteriosclerosis, Thrombosis, and Vascular Biology (2001), 21(1), 61-66  
 CODEN: ATVBFA; ISSN: 1079-5642  
 PUBLISHER: Lippincott Williams & Wilkins  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Deformation-induced endothelin-1 synthesis in endothelial cells may contribute to the intimal hyperplasia of venous bypass grafts. ACE inhibitors and angiotensin II type 1 (AT1) receptor antagonists are capable of reducing vein graft disease. Therefore, the effects of these drugs on endothelial preproendothelin-1 (ppET-1) and smooth muscle endothelin B receptor (ETB-R) expression were investigated in isolated perfused segments of the rabbit jugular vein. Pretreatment with ramiprilat (0.3 µmol/L) or irbesartan (0.01 to 1 µmol/L) had no effect on basal ppET-1 or ETB-R expression but markedly attenuated the deformation-induced expression of these gene products, and these effects were reversed by the B2 receptor antagonist icatibant (Hoe 140) and by the NO synthase inhibitor NG-nitro-L-arginine. Candesartan (1 µmol/L) mimicked the inhibitory effect of irbesartan. Moreover, reporter gene anal. with a rat ppET-1 promoter-luciferase construct transiently transfected into porcine aortic cultured endothelial cells revealed that the inhibitory effect of both ramiprilat and irbesartan on deformation-induced ppET-1 expression is species independent and mediated at the level of transcription. In addition, RT-PCR anal. detected only AT1 receptor expression in the endothelium-intact rabbit jugular vein, and neither irbesartan nor ramiprilat affected endothelial NO synthase expression. Thus, ACE inhibitors and AT1 receptor antagonists are capable of suppressing deformation-induced gene expression in the vessel wall in both an autocrine (ppET-1) and a paracrine (ETB-R) manner via a common mechanism of action that constitutes a B2 receptor-mediated increase in endothelial NO release.  
 IT 138402-11-6, Irbesartan  
 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (ACE inhibitor and AT1 antagonist blockade of deformation-induced gene expression in jugular vein through B2 receptor activation)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 31 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
 ACCESSION NUMBER: 2000:470765 CAPLUS  
 DOCUMENT NUMBER: 133:222649  
 TITLE: Synthesis and Pharmacological Evaluation of New Pyrazolidine-3,5-diones as AT1 Angiotensin II Receptor Antagonists  
 AUTHOR(S): Le'Bourdonnec, Bertrand; Meulon, Emmanuelle; Yous, Saïed; Goossens, Jean-François; Houssin, Raymond; Henichart, Jean-Pierre  
 CORPORATE SOURCE: Institut de Chimie Pharmaceutique Albert Lespagnol et Laboratoire de Chimie Analytique Faculté des Sciences Pharmaceutiques et Biologiques, Université de Lille 2, Lille, F-59006, Fr.  
 SOURCE: Journal of Medicinal Chemistry (2000), 43(14), 2685-2697  
 CODEN: JMCMAH; ISSN: 0022-2623  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB On the basis of the structure of the non-peptide receptor antagonist irbesartan, a new series of AT1 ligands was designed. In these compds. the central imidazolone nucleus of irbesartan was replaced by a pyrazolidine-3,5-dione structure. The key intermediate N-alkylpyrazolidine-3,5-diones were synthesized according to a new and general method. The most active compds. possess a spirocyclopentane ring at position 4, a linear Bu chain at position 1, and the [2'-(5-tetrazolyl)biphenyl-4-yl]methyl or [2'-(benzoylamino)sulfonyl]biphenyl-4-yl]methyl group at position 2. Affinity toward the AT1 and AT2 receptors was assessed by the ability of the compds. to competitively displace [3H]AII from its specific binding sites. The most active compds., 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-2,3-diazaspiro[4.4]nonane-1,4-dione and N-[4-[[[3-butyl-1,4-dioxo-2,3-diazaspiro[4.4]non-2-yl]methyl]-[1,1'-biphenyl]-2-yl]sulfonyl]benzamide, displayed high affinity for the AT1 receptor, good selectivity AT1 vs. AT2, and potent in vitro antagonist activity.  
 IT 138402-11-6DP, Irbesartan, derivs. and analogs  
 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (Preparation and activity of pyrazolidinediones as AT1 angiotensin II receptor antagonists)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 30 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
 REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L5 ANSWER 32 OF 36 CAPLUS COPYRIGHT 2005 ACS ON STN  
 ACCESSION NUMBER: 1999:113674 CAPLUS  
 DOCUMENT NUMBER: 130:168359  
 TITLE: Preparation of 2-oxazolinyl-4'-phthalimidomethylbiphenyls  
 INVENTOR(S): Castro, Bertrand; Dormoy, Jean-Robert; Mach, Mateusz; Makosza, Mieczyslaw; Pankowski, Jacek  
 PATENT ASSIGNER(S): Sanofi, Fr.  
 SOURCE: PCT Int. Appl., 30 pp.  
 CODEN: PIXX02  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:  

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9906398	A1	19990211	WO 1998-FR1651	19980727
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TH, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TH, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GV, ML, MR, NE, SN, TD, TG				
FR 2766821	A1	19990205	FR 1997-9653	19970729
AU 988684	A1	19990222	AU 1998-88684	19980727
PRIORITY APPL. INFO.:			FR 1997-9653	A 19970729
			WO 1998-FR1651	V 19980727
OTHER SOURCE(S):			CASREACT 130:168359; MARPAT 130:168359	
AB 4-R1C6H4C6H4(CH2R)-4 (R = phthalimido)[I] R1 = 4-(4)-(di)methyl-2-oxazolin-2-yl] were prepared as synthetic intermediates. Thus, 2-PhC6H4CO2H was cyclcondensed with Me2CH(NH2)CH2OH and the product condensed with phthalimide and trioxane to give I (R1 = 4,4-dimethyl-2-oxazolin-2-yl). The latter was used in synthesis of irbesartan a cardiovascular agent. IT 138402-10-5P RI: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (Preparation of 2-oxazolinyl-4'-phthalimidomethylbiphenyls) RN 138402-10-5 CAPLUS CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1-triphenylmethyl)-1H-tetrazol-5-yl][1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)				




REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 33 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1997:644226 CAPLUS  
 DOCUMENT NUMBER: 127:307342  
 TITLE: Radiosynthesis of [tetrazolyl-11C]irbesartan, a non-peptidic angiotensin II antagonist  
 AUTHOR(S): Ponchant, M.; Demphel, S.; Hinnen, F.; Crouzel, C.  
 CORPORATE SOURCE: DRM, Service Hospitalier Frederic-Joliot, Orsay, 91401, Fr.  
 SOURCE: European Journal of Medicinal Chemistry (1997), 32(9), 747-752  
 CODEN: EJMCAS; ISSN: 0223-5234  
 PUBLISHER: Elsevier  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB With the aim of visualizing myocardial angiotensin II receptors (AT1 subtypes), [tetrazolyl-11C]2-n-butyl-2-[(2'-(1H-tetrazol-5-yl)-1,1'-biphenyl-4-yl)methyl]-4-spirocyclopentane-2-imidazoline-5-one ([tetrazolyl-11C]irbesartan) was synthesized in one pot in four steps from [11C]hydrogen cyanide. The labeling process which yielded [tetrazolyl-11C]irbesartan is described in detail and could be applied to the labeling of other ligands which possess the (1H-tetrazol-5-yl) moiety. Positron emission tomog. (PET) studies were performed in dogs. Heart, lung and blood time-activity curves did not change. Therefore this new radioligand is not suitable for studying myocardial angiotensin II receptors with PET.  
 IT 197440-08-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (radiosynthesis of [tetrazolyl-11C]irbesartan)  
 RN 197440-08-7 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)-5-11C)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L5 ANSWER 34 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1997:519605 CAPLUS  
 DOCUMENT NUMBER: 127:185137  
 TITLE: Irbesartan. Antihypertensive, treatment of congestive heart failure, angiotensin II AT1 antagonist  
 AUTHOR(S): Casas, A.; Mellos, M.; Castaner, J.  
 CORPORATE SOURCE: Servei Nefrologia, Hospital Clinic Provincial Barcelona, Barcelona, 08080, Spain  
 SOURCE: Drugs of the Future (1997), 22(5), 481-491  
 CODEN: DRFUD4; ISSN: 0377-8282  
 PUBLISHER: Prous  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: English  
 AB A review with 66 refs. in which the authors discuss the synthesis, pharmacol. actions, clin. studies, pharmacokinetics and metabolism of irbesartan. The use of this drug as an antihypertensive, an angiotensin II AT1 antagonist and in the treatment of congestive heart failure is also discussed.  
 IT 138402-11-6P, Irbesartan  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)  
 (irbesartan synthesis, pharmacokinetics, metabolism and pharmacol. uses)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 35 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1994:671261 CAPLUS  
 DOCUMENT NUMBER: 121:271261  
 TITLE: Imidazolinones as nonpeptide angiotensin II receptor antagonists  
 AUTHOR(S): Quan, Mimi L.; DeLuca, Inda; Boswell, George A.; Chiu, Andrew T.; Wong, Pancras C.; Wexler, Ruth R.; Timmermans, Pieter B. M. W. M.  
 CORPORATE SOURCE: Dupont Merck Pharm. Co., Wilmington, DE, 19880-0402, USA  
 SOURCE: Bioorganic & Medicinal Chemistry Letters (1994), 4(12), 1527-30  
 CODEN: BMCLE8; ISSN: 0960-894X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB A series of biphenyl imidazolinones were synthesized as nonpeptide angiotensin II receptor antagonists. While those compds. with a tetrazole functionality were AT1 selective, those with a sulfonamide moiety showed affinities for both the AT1 and the AT2 sites. Representative compds. were very active in lowering blood pressure in conscious renal hypertensive rats following i.v. administration.  
 IT 138402-11-6  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (imidazolinones as nonpeptide angiotensin II receptor antagonists with antihypertensive activity)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



L5 ANSWER 36 OF 36 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1994:235405 CAPLUS  
 DOCUMENT NUMBER: 120:235405  
 TITLE: Development of tetrazole bioisosteres in angiotensin II antagonists  
 AUTHOR(S): Ferrari, B.; Taillades, J.; Perreaut, P.; Bernhart, C.; Gougat, J.; Guiraudou, P.; Cazaubon, C.; Roccon, A.; Nisato, D.; et al.  
 CORPORATE SOURCE: Sanofi Rech., Montpellier, 34184, Fr.  
 SOURCE: Bioorganic & Medicinal Chemistry Letters (1994), 4(1), 45-50  
 CODEN: BMCLE8; ISSN: 0960-894X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The application of acidic heterocycles as a substitute for tetrazole in the synthesis of potent non-peptide angiotensin II AT1 receptor antagonists is described.  
 IT 138402-11-6P  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (preparation and angiotensin II antagonist activity of)  
 RN 138402-11-6 CAPLUS  
 CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[(2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]- (9CI) (CA INDEX NAME)



```
=> s L5 and trityl?
      10788 TRITYL?
L6      4 L5 AND TRITYL?

=> d ibib abs hitstr 1-4
```


L6 ANSWER 1 OF 4 CAPIUS COPYRIGHT 2005 ACS ON STN

ACCESSION NUMBER: 2004:696369 CAPIUS

DOCUMENT NUMBER: 141:225515

TITLE: Synthesis of 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl]methyl]-1,3-diazaspiro[4.4]non-1-ene-4-one  
INVENTOR(S): Nisnevich, Gennady; Rukhman, Igor; Pertsikov, Boris; Kaftanov, Julia; Dolitzky, Ben-zion  
PATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals Usa, Inc.  
SOURCE: PCT Int. Appl., 27 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004072064	A1	20040826	WO 2004-US3604	20040205
V: AE, AE, AG, AL, AL, AM, AM, AM, AT, AT, AU, AU, AZ, AZ, BA, BA, BG, BG, BR, BR, BV, BY, BY, BZ, BZ, CA, CA, CH, CN, CN, CO, CO, CR, CR, CU, CU, CZ, CZ, DE, DE, DK, DK, DM, DM, EC, EC, EE, EE, EG, EG, ES, ES, FI, FI, GB, GB, GE, GE, GH, GH, GR, GR, HU, HU, ID, ID, IL, IL, IS, JP, JP, KE, KE, KG, KG, KP, KP, KR, KR, KZ, KZ, LC, LC, LK, LR, LS, LS, LT, LU, LV, MA, MD, MD, MG, MK, MN, MW, MX, MX, MZ, MZ, NA, NI				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZV, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2004242894	A1	20041202	US 2004-773414	20040205
PRIORITY APPL. INFO.:			US 2003-445218P	P 20030205
			US 2003-465905P	P 20030428
OTHER SOURCE(S):	CASREACT 141:225515			
GI				



I

AB Provided are 5 methods of making 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl]methyl]-1,3-diazaspiro[4.4]non-1-ene-4-one (I), e.g. comprising the steps of: (a) reacting 1-(N'-pentanoylamino)cyclopentanecarboxylic acid amide with 5-(4'-

L6 ANSWER 2 OF 4 CAPIUS COPYRIGHT 2005 ACS ON STN

ACCESSION NUMBER: 2004:633925 CAPIUS

DOCUMENT NUMBER: 141:157121

TITLE: Synthesis of irbesartan  
INVENTOR(S): Nisnevich, Gennady; Rukhman, Igor; Pertsikov, Boris; Kaftanov, Julia; Dolitzky, Ben-zion  
PATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals Usa, Inc.  
SOURCE: PCT Int. Appl., 21 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004065383	A2	20040805	WO 2004-US1135	20040116
WO 2004065383	A3	20041216		
V: AE, AE, AG, AL, AL, AM, AM, AM, AT, AT, AU, AU, AZ, AZ, BA, BA, BG, BG, BR, BR, BV, BY, BY, BZ, BZ, CA, CA, CH, CN, CN, CO, CO, CR, CR, CU, CU, CZ, CZ, DE, DE, DK, DK, DM, DM, EC, EC, EE, EE, EG, EG, ES, ES, FI, FI, GB, GB, GE, GE, GH, GH, GR, GR, HU, HU, ID, ID, IL, IL, IS, JP, JP, KE, KE, KG, KG, KP, KP, KR, KR, KZ, KZ, LC, LC, LK, LR, LS, LS, LT, LU, LV, MA, MD, MD, MG, MK, MN, MW, MX, MX, MZ				
US 2004192713	A1	20040930	US 2004-759906	20040116
EP 1509517	A2	20050302	EP 2004-702955	20040116
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
PRIORITY APPL. INFO.:			US 2003-440997P	P 20030116
			WO 2004-US1135	W 20040116

OTHER SOURCE(S): CASREACT 141:157121


AB Provided are a method of making irbesartan via a Suzuki coupling reaction and a novel intermediate, 2-butyl-3-(4'-bromobenzyl)-1,3-diazaspiro[4.4]non-1-ene-4-one, for such process. The novel process includes the step of reacting such intermediate with a protected tetrazolylphenylboronic acid.

IT 138402-11-6P, Irbesartan  
RL: INF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)

(synthesis of irbesartan)

RN 138402-11-6 CAPIUS

CN 1,3-Diazaspiro[4.4]non-1-ene-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



IT 138402-10-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(synthesis of irbesartan)

L6 ANSWER 1 OF 4 CAPIUS COPYRIGHT 2005 ACS ON STN (Continued)


bromomethylbiphenyl-2-yl)-1-trityl-1H-tetrazole in the presence of an inorg. base, a solvent and a phase transfer catalyst; (b) cooling the mixt.; (c) adding water to the mixt. whereby two phases are obtained; (d) sepg. the two phases obtained; and (e) recovering the compd. I. The compds. I can be converted to irbesartan which is a known angiotensin II receptor antagonist (blocker).

IT 745814-09-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(methods for preparation of 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl]methyl]-1,3-diazaspiro[4.4]non-ene-4-one)

RN 745814-09-9 CAPIUS

CN 1,3-Diazaspiro[4.4]non-1-ene-4-one, 2-[[1,1-dimethylethyl]-3-[[2'-(1-triphenylmethyl)-1H-tetrazol-5-yl][1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




IT 138402-11-6P, Irbesartan

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(methods for preparation of 2-butyl-3-[[2'-(1-trityl-1H-tetrazol-5-yl)biphenyl-4-yl]methyl]-1,3-diazaspiro[4.4]non-ene-4-one)

RN 138402-11-6 CAPIUS


CN 1,3-Diazaspiro[4.4]non-1-ene-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



L6 ANSWER 2 OF 4 CAPIUS COPYRIGHT 2005 ACS ON STN (Continued)

RN 138402-10-5 CAPIUS

CN 1,3-Diazaspiro[4.4]non-1-ene-4-one, 2-butyl-3-[[2'-(1-triphenylmethyl)-1H-tetrazol-5-yl][1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



*Copyright*

ACCESSION NUMBER: 2004:50495 CAPLUS

DOCUMENT NUMBER: 140:111420

TITLE:

INVENTOR(S): Synthesis of irbesartan  
Nisnevich, Gennady; Rukhman, Igor; Pertsikov, Boris;  
Kaftanov, Julia; Dolitzky, Ben-Zion; Shapiro, Eugeny;  
Yahalom, BonitPATENT ASSIGNEE(S): Teva Pharmaceutical Industries Ltd., Israel; Teva  
Pharmaceuticals USA, Inc.

SOURCE: PCT Int. Appl., 13 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007482	A2	20040122	WO 2003-US22479	20030716
WO 2004007482	A3	20040527		
V:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RV:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2492779	AA	20040122	CA 2003-2492779	20030716
EP 1546135	A2	20050629	EP 2003-764805	20030716
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
US 2005176794	A1	20050911	US 2003-621623	20030716
PRIORITY APPLN. INFO.:			US 2002-396424P	P 20020716
			US 2002-402490P	P 20020809
			WO 2003-US22479	W 20030716

OTHER SOURCE(S): CASREACT 140:111420

AB Irbesartan is prepared by reaction of 2-butyl-1,3-diazaspiro[4.4]non-1-ene (I) with 5-(4-bromomethylbiphenyl-2-yl)-1-trityl-1H-tetrazole (II) in the presence of a phase transfer catalyst. Thus, reaction of I with II in toluene in the presence of Bu<sub>4</sub>NHSO<sub>4</sub> at 90° for 1.5 h gave, after deprotection, 84.3% irbesartan. Also provided is irbesartan having a fine particle size.


IT 138402-11-6P, Irbesartan

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of irbesartan)

RN 138402-11-6 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)

(Continued)




IT 138402-10-5

RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of irbesartan)

RN 138402-10-5 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1-(triphenylmethyl)-1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



Copyright

ACCESSION NUMBER: 1999:113674 CAPLUS

DOCUMENT NUMBER: 130:168359

TITLE:

INVENTOR(S): Preparation of 2-oxazoliny-4'-  
phthalimidomethylbiphenyls  
Castro, Bertrand; Dormoy, Jean-Robert; Mach, Mateusz;  
Makosza, Mieczyslaw; Pankowski, Jacek

PATENT ASSIGNEE(S): Sanoel, F.

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9906398	A1	19990211	WO 1998-FR1651	19980727
V:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RV:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, BG, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2766821	A1	19990205	FR 1997-9653	19970729
AU 9888684	A1	19990222	AU 1998-88684	19980727
PRIORITY APPLN. INFO.:			FR 1997-9653	A 19970729
			WO 1998-FR1651	W 19980727


OTHER SOURCE(S): CASREACT 130:168359; MARPAT 130:168359  
AB 4-RICG14CGH4 (CH<sub>2</sub>R)-4 (R = phthalimido)[1,1'-4(4)-(di)methyl-2-oxazolin-2-yl] were prepared as synthetic intermediates. Thus, 2-PHC6H4CO<sub>2</sub>H was cyclocondensed with Me<sub>2</sub>CH(NH<sub>2</sub>)CH<sub>2</sub>OH and the product condensed with phthalimide and trioxane to give I (R<sub>1</sub> = 4,4-dimethyl-2-oxazolin-2-yl). The latter was used in synthesis of irbesartan a cardiovascular agent.

IT 138402-10-5P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of 2-oxazoliny-4'-phthalimidomethylbiphenyls)

RN 138402-10-5 CAPLUS

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1-(triphenylmethyl)-1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)




IT 138402-11-6P, Irbesartan

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of 2-oxazoliny-4'-phthalimidomethylbiphenyls)

RN 138402-11-6 CAPLUS

(Continued)

CN 1,3-Diazaspiro[4.4]non-1-en-4-one, 2-butyl-3-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 7

THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT